

# ARIZONA GUIDE TO IMMUNIZATIONS REQUIRED FOR ENTRY

## Grades K-12 (School year 2022-2023)

- Requirements are shown below as stated in [Arizona Administrative Code, R9-6-702](#), Table 7.1 and Table 7.2
- Please review the [Arizona Immunization Handbook for Schools and Child Care Programs](#) along with the [Vaccine Catch-up Flowcharts & FAQs](#) for further information and details about immunization requirements and exemptions.
- Vaccines must follow ACIP minimum intervals and ages to be valid. See page 2 for minimum interval and catch-up schedule information.
- The 4-day grace period only applies to vaccine-administration minimum age and intervals. Refer to the Handbook for questions.



Vaccine	4-6 Years Old and attendance in Kindergarten or 1 <sup>st</sup> grade	7-10 Years Old	11 Years and Older
<b>HepB</b> Hepatitis B	<b>3 doses</b> The final dose of HepB must be given at 24 weeks of age or older. Only 3 doses are required if the 3 <sup>rd</sup> dose was received at or after the child was 24 weeks of age; otherwise 4 doses are required.		
<b>Polio</b> Poliomyelitis (IPV) For OPV see page 2	<b>4 doses</b> The final dose of polio must be received at/after 4 years of age and at least six months after the previous dose. Only 3 doses are required if the 3 <sup>rd</sup> dose was received on/after the child's 4 <sup>th</sup> birthday and at least six months after the 2 <sup>nd</sup> dose. Additional doses may be needed to meet requirements. See pg. 2 for retrospective history guidance.		
<b>MMR</b> Measles, Mumps and Rubella	<b>2 doses</b> Minimum recommended age for dose 1 is 12 months. A 3 <sup>rd</sup> dose will be required if dose 1 was given more than 4 days before 1 <sup>st</sup> birthday.		
<b>VAR</b> Varicella (chickenpox)	<b>1 dose</b> Minimum recommended age for dose 1 is 12 months. 2 doses, at least 4 weeks apart, are required if dose 1 was given at 13 years of age or older.		
<b>DTaP, Tdap, Td</b> Diphtheria, Tetanus, and Pertussis	<b>5 doses of DTaP</b> The final dose of tetanus-diphtheria containing vaccine must be received at/after 4 years of age and at least six months after the previous dose. Only 4 doses are required if the 4 <sup>th</sup> dose was received on/after 4 years of age; in certain situations, an additional dose may be required, up to a maximum of 6 doses (before age 7).	<b>4 doses of tetanus-diphtheria-containing vaccine (or combination of DTaP, Td or Tdap doses). At least one dose at/after 4 years of age and at least 6 months from previous dose.</b>  3 doses (with one at/after 4 years) is acceptable if the first dose was given on/after 1 <sup>st</sup> birthday; otherwise refer for an additional dose.	<b>1 dose of Tdap is required</b> If the student does not have a Tdap but received a dose of tetanus-diphtheria-containing vaccine within the past 5 years, refer for the adolescent Tdap dose when 5 years has passed since that dose. If a student has received 1 valid dose of adolescent Tdap (age 10 years or older), no further doses are needed.  Students must have minimum series of 4 doses of tetanus-diphtheria-containing vaccine; 3 doses acceptable if the 1 <sup>st</sup> dose was given on/after 1 <sup>st</sup> birthday.
<b>MenACWY or MCV4</b> Quadrivalent Meningococcal	Retrospectively: Menomune (Meningococcal Polysaccharide) vaccine was a quadrivalent vaccine so is acceptable; however, production of this vaccine was discontinued in February 2017. Menomune doses are considered acceptable for school requirements.		<b>1 dose of MenACWY is required</b> A dose administered at 10 years of age will meet the requirement.

## ARIZONA GUIDE TO IMMUNIZATIONS REQUIRED FOR ENTRY – Minimum Interval/Catch-up Guidance

### Grades K-12 (School year 2022-2023)

Vaccine	Dose #	Minimum Age	Minimum Interval Between Doses	Notes
<b>HepB</b> Hepatitis B	dose 1	Birth	At least 4 weeks between dose 1 & 2	<ul style="list-style-type: none"> <li>Some children may receive a birth dose and then a combination vaccine resulting in a total of 4 (or more) doses. At long as the interval between doses is met, 4+ doses meet requirement.</li> <li>2 doses, at least 4 months apart, meet the requirement if the child received the adolescent series using the Merck Recombivax HB Adult Formulation when the child was 11-15 years of age.</li> </ul>
	dose 2	4 weeks	At least 8 weeks between dose 2 & 3 (or final)	
	dose 3	24 weeks	At least 16 weeks between dose 1 & 3 (or final) AND at/after 24 weeks of age	
<b>Polio</b> IPV or OPV	dose 1	6 weeks	At least 4 weeks between dose 1 & 2	<ul style="list-style-type: none"> <li>Retrospectively: 1) A final dose given on or after August 7, 2009, must be given at or after 4 years of age <b>and</b> a minimum interval of 6 months from the previous dose. 2) Students who received 4 doses (with at least 4 weeks minimum intervals between doses and/or before the age of 4 years) PRIOR to August 7, 2009 have met the requirement.</li> <li>OPV given prior to April 1, 2016 will be presumed to be trivalent and therefore acceptable, regardless of age, or country, of administration. Any OPV doses administered on or after April 1, 2016 are presumed to be bivalent and therefore unacceptable.</li> <li>Poliomyelitis vaccine is not recommended in the U.S. for individuals 18 years of age or older; however, a complete series is still required for school attendance.</li> </ul>
	dose 2	10 weeks	At least 4 weeks between dose 2 & 3	
	dose 3	14 weeks	At least 4 weeks between dose 3 & 4	
	dose 4	4 years	At least 6 months between final dose and previous dose (could be final dose 3 or final dose 4)	
<b>MMR</b> Measles, Mumps and Rubella	dose 1	12 months	At least 4 weeks (28 days) between dose 1 & 2	<ul style="list-style-type: none"> <li>If MMR dose 1 was given more than 4 days before the 1<sup>st</sup> birthday, another dose is required.</li> <li>MMR and varicella vaccines are live vaccines and must be given on the same day or at least 28 days apart (this rule also applies to live nasal influenza doses).</li> </ul>
	dose 2	13 months	-	
<b>VAR</b> Varicella (chickenpox)	dose 1	12 months	At least 3 months between dose 1 & 2  4 weeks (28 days) between doses if administered at age 13 or older	<ul style="list-style-type: none"> <li>If varicella dose 1 was given more than 4 days before the 1<sup>st</sup> birthday, another dose is required.</li> <li>MMR and varicella vaccines are live vaccines and must be given on the same day or at least 28 days apart (this rule also applies to live nasal influenza doses).</li> </ul>
<b>DTaP, Tdap, Td</b> Tetanus, Diphtheria, and Pertussis	dose 1	6 weeks	At least 4 weeks between dose 1 & 2	<ul style="list-style-type: none"> <li>DTaP is licensed for children through age 6. If catch-up doses are needed at age 7 or older, Tdap or Td should be used to start/complete the series.</li> <li>A Tdap given at age 7-9 years of age does not count for the 11- year old Tdap requirement; a Tdap should be given once 5 years has passed since last dose of tetanus-diphtheria containing vaccines was given.</li> <li>Retrospectively, if a child received a Tdap at age 10 as part of a catch-up series, or inadvertently earlier than the recommended age of 11-12, the dose may be counted as the adolescent dose and is acceptable to meet school requirements.</li> <li>Once a valid adolescent Tdap dose has been received, a tetanus booster is recommended when 10 years has passed since last dose of tetanus-containing vaccine.</li> </ul>
	dose 2	10 weeks	At least 4 weeks between dose 2 & 3	
	dose 3	14 weeks	At least 6 months between dose 3 & 4	
	dose 4	12 months	At least 6 months between dose 4 & 5	
	dose 5	4 years	In general, a child should not receive more than 4 doses prior to the 4 <sup>th</sup> birthday or a total of 6 doses prior to the 7th birthday; however, the child should still receive a dose at/after 4 years of age and at least 6 months from previous dose	
<b>MenACWY, MCV4</b> Meningococcal	dose 1	10 years		<ul style="list-style-type: none"> <li>Only quadrivalent meningococcal ACWY vaccine doses will be accepted. The vaccines given currently in the U.S. are Menactra, Menveo, and MenQuadfi. No monovalent or bivalent meningococcal vaccinations will be accepted (MenA, MenB, MenC, or MenC/Y).</li> </ul>